

▼ This medicinal product is subject to additional monitoring in Australia. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse events at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

## AUSTRALIAN PRODUCT INFORMATION

### **XEVUDY (Sotrovimab) Concentrated injection solution for infusion**

#### **1 NAME OF THE MEDICINE**

Sotrovimab

#### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each vial contains 500 mg of sotrovimab in 8 mL (62.5 mg/mL).

For the full list of excipients, see Section 6.1 LIST OF EXCIPIENTS.

#### **3 PHARMACEUTICAL FORM**

Sotrovimab is a clear, colourless or yellow to brown concentrated injection solution for intravenous infusion.

#### **4 CLINICAL PARTICULARS**

##### **4.1 THERAPEUTIC INDICATIONS**

XEVUDY has **provisional approval** for the treatment of adults and adolescents (aged 12 years and over and weighing at least 40 kg) with coronavirus disease 2019 (COVID-19) who do not require initiation of oxygen due to COVID-19 and who are at increased risk of progression to hospitalisation or death (see Section 5.1 PHARMACODYNAMIC PROPERTIES, Clinical Trials).

The decision has been made on the basis of short term efficacy and safety data. Continued approval of this indication depends on the evidence of longer term efficacy and safety from ongoing clinical trials and post-market assessment.

##### **4.2 DOSE AND METHOD OF ADMINISTRATION**

As part of risk stratification of patients the pivotal consideration is the comorbidities, alongside age, particularly multiple comorbidities.

XEVUDY should not be used in patients hospitalised due to COVID-19.

##### **Adults and adolescents (aged 12 years and older and weighing at least 40 kg)**

The recommended regimen is a single 500 mg dose administered as an intravenous infusion.

##### **Method of Administration**

XEVUDY is administered as a single intravenous (IV) infusion over 30 minutes (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE).

It is recommended that XEVUDY is administered within 5 days of onset of symptoms of COVID-19 (see Section 5.1 PHARMACODYNAMIC PROPERTIES, Clinical Trials).

XEVUDY must be diluted prior to administration and must not be administered as an intravenous push or bolus injection.

XEVUDY should be administered in healthcare facilities in which patients can be monitored during and for one hour after administration of XEVUDY (see Section 4.4. SPECIAL WARNINGS AND PRECAUTIONS FOR USE).

#### Preparation for Dilution

XEVUDY should be prepared by a qualified healthcare professional using aseptic technique.

1. Remove one vial of XEVUDY from the refrigerator (2°C to 8°C). Allow the vial to equilibrate to ambient room temperature, protected from light, for approximately 15 minutes.
2. Visually inspect the vial to ensure it is free from particulate matter and that there is no visible damage to the vial.
  - a. If a vial is identified to be unusable, discard and restart the preparation with a new vial.
3. Gently swirl the vial several times before use without creating air bubbles.
  - a. Do not shake or vigorously agitate the vial.

#### Dilution Instructions for Intravenous Infusion

1. Withdraw 8 mL from an infusion bag containing 50 mL or 100 mL of sodium chloride 9mg/mL (0.9%) solution for injection or 5% dextrose for injection.
2. Withdraw 8 mL from the vial of XEVUDY.
3. Inject the 8 mL of XEVUDY into the infusion bag via the septum.
4. Discard any unused portion left in the vial as the product contains no preservative. The vial is single-use only and should only be used for one patient.
5. Prior to the infusion, gently rock the infusion bag back and forth 3 to 5 times. Do not invert the infusion bag. Avoid forming air bubbles.

The diluted solution of XEVUDY is intended to be used immediately. If immediate administration is not possible, the diluted solution may be stored at room temperature (up to 25°C) for up to 6 hours or refrigerated (2°C to 8°C) for up to 24 hours from the time of dilution until the end of administration.

#### Administration Instructions

1. Attach an infusion set to the infusion bag using standard bore tubing. The intravenous dosing solution is recommended to be administered with a 0.2-µm in-line filter.
2. Prime the infusion set.
3. Administer as an IV infusion over 30 minutes at room temperature.

## **Children**

The safety and efficacy of XEVUDY have not been established in children less than 12 years of age or weighing less than 40 kg (see Section 5.2 PHARMACOKINETIC PROPERTIES, Special patient populations).

## **Elderly**

No dose adjustment is required in patients aged 65 years or older (see Section 5.2 PHARMACOKINETIC PROPERTIES, Special patient populations).

## **Renal impairment**

No dose adjustment is required in patients with renal impairment (see Section 5.2 PHARMACOKINETIC PROPERTIES, Special patient populations).

## **Hepatic impairment**

No dose adjustment is required in patients with hepatic impairment (see Section 5.2 PHARMACOKINETIC PROPERTIES, Special patient populations).

## **4.3 CONTRAINDICATIONS**

Hypersensitivity to the active substance or any of the excipients (see Section 6.1 LIST OF EXCIPIENTS).

## **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE**

### **Hypersensitivity reactions**

In a study in patients hospitalised with COVID-19, hypersensitivity reactions, including serious reactions such as anaphylaxis, have been reported following infusion of sotrovimab. If signs and symptoms of a clinically significant hypersensitivity reaction occur, immediately discontinue administration and initiate appropriate supportive care.

In COMET-ICE, mild to moderate hypersensitivity reactions have been observed. If mild to moderate hypersensitivity reactions occur, consider slowing or stopping the infusion along with appropriate supportive care.

### **Use in elderly**

Based on population pharmacokinetic analysis, there was no difference in sotrovimab pharmacokinetics in elderly patients when compared with younger patients.

### **Paediatric use**

The safety and efficacy of XEVUDY have not been established in children less than 12 years of age or weighing less than 40 kg (see Section 5.2 PHARMACOKINETIC PROPERTIES, Special patient populations).

### **Effects on laboratory tests**

No data available.

## **4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS**

No formal interaction studies have been conducted with sotrovimab.

Sotrovimab is not renally excreted or metabolized by cytochrome P450 (CYP) enzymes; therefore, interactions with concomitant medications that are renally excreted or that are substrates, inducers, or inhibitors of CYP enzymes are unlikely.

In *in vitro* pharmacodynamic studies with remdesivir or bamlanivimab, sotrovimab showed additive virologic effect and no antagonism with either agent.

The efficacy and safety of sotrovimab in subjects who have received a COVID-19 vaccine at any time prior to its administration has not been established. The receipt of a COVID-19 vaccine within 48 hours prior to, or 4 weeks following treatment with sotrovimab has not been studied.

## **4.6 FERTILITY, PREGNANCY AND LACTATION**

### **Effects on fertility**

There are no data on the effects of sotrovimab on human male or female fertility. Effects on male and female fertility have not been evaluated in animal studies.

### **Use in pregnancy (Category B2)**

There are insufficient data on the effects of sotrovimab on human pregnancy. Effects on embryo-fetal development have not been evaluated in animal studies. In a cross-reactive binding assay using a protein array enriched for human embryofetal proteins, no off-target binding was detected. Since sotrovimab is an engineered human immunoglobulin G (IgG), it has the potential for placental transfer from the mother to the developing fetus. The potential treatment benefit or risk of placental transfer of sotrovimab to the developing fetus is not known.

XEVUDY should be used during pregnancy only if the expected benefit to the mother justifies the potential risk to the foetus.

### **Use in lactation**

There are insufficient data on the presence of sotrovimab in human milk. There are no data in lactating animals. A decision must be made whether to discontinue breast-feeding or to abstain from sotrovimab therapy considering the benefit of breast-feeding for the child and the benefit of therapy for the mother.

## **4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES**

There have been no studies to investigate the effect of sotrovimab on the ability to perform tasks that require judgement, motor or cognitive skills. A detrimental effect on such activities would not be anticipated from the pharmacology of sotrovimab. The clinical status of the patient and the adverse event profile of sotrovimab should be borne in mind when considering the patient's ability to perform tasks that require judgement, motor or cognitive skills.

## 4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)

### Clinical trial data

The safety of sotrovimab was evaluated in a placebo-controlled randomised study in 1049 non-hospitalised patients with COVID-19 (COMET-ICE) (see Section 5.1 PHARMACODYNAMIC PROPERTIES, Clinical trials).

**Table 1: Summary of common adverse events reported in the COMET-ICE trial at an incidence of  $\geq 1\%$ .**

MedDRA System Organ Class Preferred Term	Sotrovimab 500 mg (n = 526) n (%)	Placebo (n = 523) n (%)
Gastrointestinal disorders		
Nausea	5 (<1%)	9 (2%)
Diarrhea	8 (2%)	4 (<1%)
Infections and infestations		
COVID-19 pneumonia <sup>a</sup>	5 (<1%)	22 (4%)
Nervous system disorders		
Headache	4 (<1%)	11 (2%)

<sup>a</sup>As recorded by the investigator

Adverse reactions are listed below by MedDRA body system organ class (SOC) and by frequency (Table 2). Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  and  $< 1/10$ ), uncommon ( $\geq 1/1,000$  and  $< 1/100$ ), rare ( $\geq 1/10,000$  and  $< 1/1,000$ ) and very rare ( $< 1/10,000$ ).

**Table 2: Adverse reactions**

System Organ Class	Frequency	Adverse Reactions
Immune system disorders	Common	Hypersensitivity reactions <sup>a</sup>

<sup>a</sup>Includes rash, dermatitis contact, skin reaction, hypersensitivity, multiple allergies, infusion-related reaction and bronchospasm

### Hypersensitivity including anaphylaxis and infusion-related reactions

In COMET-ICE, hypersensitivity reactions, of grade 1 (mild) or grade 2 (moderate), were reported (9 patients in the sotrovimab arm; 5 patients in the placebo arm). None of the reactions in either arm led to pausing or discontinuation of the infusions.

One case of anaphylaxis was reported following infusion of sotrovimab in a study in hospitalized patients; the patient received adrenaline (epinephrine) and the event resolved.

### Post-marketing data

There are no relevant data available.

### Reporting suspected adverse effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product.

Healthcare professionals are asked to report any suspected adverse reactions at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

## 4.9 OVERDOSE

There is no clinical experience with overdose of sotrovimab.

There is no specific treatment for an overdose of sotrovimab. If overdose occurs, the patient should be treated supportively with appropriate monitoring as necessary.

For information on the management of overdose, contact the Poisons Information Centre on 13 11 26 (Australia).

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 PHARMACODYNAMIC PROPERTIES

Pharmacotherapeutic group: immune sera and immunoglobulins: immunoglobulins, specific immunoglobulins.

#### ATC code

Not yet assigned

#### **Mechanism of action**

Sotrovimab is an engineered human IgG1 mAb that binds to a highly conserved epitope on the spike protein receptor binding domain of SARS-CoV-2 with high affinity (dissociation constant  $K_d = 31$  ng/mL). The Fc domain of sotrovimab includes M428L and N434S amino acid substitutions (LS modification) that extends antibody elimination half-life, but does not impact wild-type Fc-mediated effector functions when compared with the original mAb with LS modification (S309+LS) in cell culture

#### **Pharmacodynamic effects**

##### Immunogenicity

Consistent with the potentially immunogenic properties of protein and peptide therapeutics, patients may develop antibodies to sotrovimab following treatment. The clinical relevance of such antibodies has not been fully established.

##### Antiviral activity

Sotrovimab neutralized SARS-CoV-2 *in vitro* (76.6 – 132.5 ng/mL), and *in vivo* ( $\geq 5$  mg/kg in SARS-CoV-2 infected hamsters dosed with sotrovimab prior to virus inoculation) and effectively neutralised pseudotyped virus containing the SARS-CoV-2 spike.

Sotrovimab exhibited antibody-dependent cell-mediated cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP) in cell-based assays.

Sotrovimab demonstrated activity *in vivo* in a hamster model of SARS-CoV-2 infection using sotrovimab as well as VIR-7831-wild type (WT), a mAb that has identical variable regions as sotrovimab but is lacking the LS modification. Intraperitoneal administration of sotrovimab or VIR-7831-WT at  $\geq 5$  mg/kg prior to inoculation resulted in a significant improvement in body

weight loss. Sotrovimab and VIR-7831-WT significantly decreased total viral RNA in the lungs at  $\geq 0.5$  and  $\geq 5$  mg/kg, respectively, and infectious virus levels based on TCID<sub>50</sub> measurements at  $\geq 0.5$  mg/kg.

### Antibody Dependent Enhancement (ADE)

The risk that sotrovimab could mediate viral uptake and replication by immune cells was studied in U937 and primary human monocytic dendritic cells and peripheral blood mononuclear cells. This experiment did not demonstrate productive viral infection in immune cells exposed to SARS-CoV-2 at concentrations of sotrovimab from 1-fold down to 1000-fold the EC<sub>50</sub> value.

The potential for ADE was also evaluated in a hamster model of SARS-CoV-2 using sotrovimab as well as VIR-7831-wild type (WT). No evidence of enhancement of disease was observed at any dose evaluated, including sub-neutralizing doses down to 0.05 mg/kg. Additionally, a separate hamster study using a modified version of the parental antibody S309 that interacts with hamster FcRs was conducted. There was no evidence of ADE using the modified antibody at neutralising or sub-neutralising doses.

### Antiviral Resistance

There is a potential risk of treatment failure due to the development of viral variants that are resistant to sotrovimab.

A pseudotyped VLP assessment in cell culture and/or *in vitro* resistance selection with increasing concentrations of sotrovimab showed that the epitope sequence polymorphisms at P337 and E340 confer reduced susceptibility to sotrovimab. EC<sub>50</sub> values against P337H/L/R/T and E340A/K/G increased by 5.4- $\rightarrow$ 297 fold relative to the wild type. Pseudotyped VLP *in vitro* assessments indicate that sotrovimab retains activity against the UK (Alpha, B.1.1.7; 2.30-fold change in EC<sub>50</sub> value); South Africa (Beta, B.1.351; 0.60-fold change in EC<sub>50</sub> value); Brazil (Gamma, P.1; 0.35-fold change in EC<sub>50</sub> value); California (Epsilon, B.1.427/B.1.429; 0.70-fold change in EC<sub>50</sub> value); New York (Iota, B.1.526; 0.6-fold change in EC<sub>50</sub> value) and India (Kappa, B.1.617; 0.7-fold change in EC<sub>50</sub> value, Delta, B.1.617.2; 1-fold change in EC<sub>50</sub> value) variant spike proteins. Microneutralisation data from authentic SARS-CoV-2 variant virus also indicate that sotrovimab retains activity against the UK (3-fold change in EC<sub>50</sub> value), South Africa (1.2-fold change in EC<sub>50</sub> value) and Brazil (1.6-fold change in EC<sub>50</sub> value) variants.

In the COMET-ICE clinical trial, based on sequences of 45 participant in the sotrovimab arm with available post-baseline data, post-baseline epitope variants were detected in 10 participants in the sotrovimab arm (A344V [6.2%]; R346G [5.2%]; K356R [7.5%]; 2 patients with S359G [12.2% and 8.3%]; C361T [6.0%]; 4 patients with E340K [ $\geq 99.7\%$ ]). Of the variants detected at baseline and post-baseline, L335F, L335S, G339C, E340A, E340K, A344V, R346I, R346G, K356N, K356R, R357I, I358V and S359G have been assessed phenotypically using a pseudotyped VLP system. Sotrovimab retains susceptibility against L335F (0.8-fold change in EC<sub>50</sub> value), L335S (0.9-fold change in EC<sub>50</sub> value), G339C (1.2-fold change in EC<sub>50</sub> value), A344V (1.1-fold change in EC<sub>50</sub> value), R346I (1.7-fold change in EC<sub>50</sub> value), R346G (0.9 change in EC<sub>50</sub> value), K356N (1.1-fold change in EC<sub>50</sub> value), K356R (0.8-fold change in EC<sub>50</sub> value), R257I (1-fold change in EC<sub>50</sub> value), I358V (0.7-fold change in EC<sub>50</sub> value), and S359G (0.8-fold change in EC<sub>50</sub> value). E340A and E340K

confer reduced susceptibility to sotrovimab (>100-fold and >297-fold change in EC<sub>50</sub> value, respectively).

The clinical impact of these variants is not yet known. Data collection and analysis is still ongoing.

### **Clinical trials**

Study 214367 (COMET-ICE) was a Phase II/III randomised, double-blind, placebo-controlled study which evaluated sotrovimab as treatment for COVID-19 in non-hospitalised patients at high risk of medical complications of the disease. Patients included were aged 18 years and older with at least 1 of the following comorbidities: diabetes requiring medication, obesity (BMI>30), chronic kidney disease (eGFR< 60 mL/min), congestive heart failure (NYHA ≥ class 2), chronic obstructive pulmonary disease, or moderate to severe asthma (requiring inhaled steroids to control the symptoms or has been prescribed a course of oral corticosteroids in the past year), or were aged 55 years and older. The study included symptomatic patients with SARS-CoV-2 infection, as confirmed by local laboratory tests and/or point of care tests. Patients with severe COVID-19 requiring supplemental oxygen or hospitalization and patients who have received a COVID-19 vaccine were excluded from the trial. Patients were randomised to receive a single 500 mg infusion of sotrovimab (N = 528) or placebo (N = 529) over 1 hour (Intent to Treat [ITT] population at Day 29).

A total of 46% of randomised participants were male. The median age of the overall randomised population was 53 years (range: 17 to 96). A total of 20% of participants were aged 65 years or older and 11% were over 70 years of age. The majority of participants were of White race (87%); 8% were Black or African American and 4% were Asian. The ethnicity of the majority of participants was Hispanic or Latino (65%). 59% of participants received the sotrovimab or placebo within 3 days of COVID-19 symptom onset and 41% within 4-5 days. The four most common pre-defined risk factors or comorbidities were obesity (63%), 55 years of age or older (47%), diabetes requiring medication (22%) and moderate to severe asthma (17%). Overall, baseline demographic and disease characteristics were well balanced between the treatment arms.

Enrollment to COMET-ICE was halted for overwhelming efficacy following a pre-specified interim analysis (IA) of the primary endpoint. The primary endpoint, progression of COVID-19 at Day 29, was reduced by 79% compared with placebo (adjusted relative risk reduction) in recipients of sotrovimab (p<0.001) in the final intention to treat (ITT) population. This reduction is consistent in magnitude to that previously reported for ITT (IA) population. Tables 3 and 4 below, provides the results of the primary endpoint and key secondary endpoints of COMET-ICE.

**Table 3: Results of primary and secondary endpoints in the ITT population at Day 29 (COMET-ICE)**

	<b>Sotrovimab (500 mg IV infusion) N= 528</b>	<b>Placebo N= 529</b>
<b>Primary endpoint</b>		
<b>Progression of COVID-19 as defined by hospitalisation for &gt;24 hours for acute management of any illness or death from any cause (Day 29)</b>		
Proportion (n, %) <sup>a</sup>	6 (1%)	30 (6%)
Adjusted relative risk reduction (95% CI)	79% (50%, 91%)	
p-value	<0.001	
<b>Secondary endpoints</b>		
<b>Progression of COVID-19 as defined by visit to a hospital emergency room for management of illness or hospitalisation for acute management of illness or death from any cause (Day 29)</b>		
Proportion (n, %)	13 (2%)	39 (7%)
Adjusted relative risk reduction (95% CI)	66% (37%, 81%)	
p-value	<0.001	
<b>Progression to develop Severe and/or Critical Respiratory COVID-19 (Day 29) <sup>b</sup></b>		
Proportion (n, %) <sup>c</sup>	7 (1%)	28 (5%)
Adjusted relative risk reduction (95% CI)	74% (41%, 88%)	
p-value	0.002	
<b>All-cause mortality (up to Day 29)</b>		
Proportion (n, %)	0	2 (<1%)
<sup>a</sup> No participants required intensive care unit (ICU) stay in the sotrovimab arm versus 9 participants in the placebo arm. <sup>b</sup> Progression to develop severe and/or critical respiratory COVID-19 defined as the requirement for supplemental oxygen (low flow nasal cannulae/face mask, high flow oxygen, non-invasive ventilation, mechanical ventilation or extracorporeal membrane oxygenation [ECMO]). <sup>c</sup> No participants required use of high flow oxygen, non-rebreather mask or mechanical ventilation in the sotrovimab arm versus 14 participants in the placebo arm.		

**Table 4: Summary of Nasal SARS-CoV-2 Viral Load in log 10 copies/mL on Day 8 in the Virology Population (secondary endpoint)**

	Sotrovimab (500 mg IV infusion)	Placebo
<b>Baseline (log 10 copies/mL)</b>		
n	358	375
Mean (standard deviation)	6.554 (1.6248)	6.652 (1.6732)
<b>Day 8 (log 10 copies/mL)</b>		
n <sup>a</sup>	294	305
Mean (standard error)	4.005 (0.0606)	4.236 (0.0598)
<b>Day 8 change from baseline (log 10 copies/mL)</b>		
Mean (standard error)	-2.589 (0.0606)	-2.357 (0.0598)
95% CI	-2.708, -2.470	-2.475, -2.240
Difference (standard error)	-0.232 (0.0851)	
95% CI	-0.399, -0.065	
p-value	0.007	

<sup>a</sup> Number of participants with available data at Day 8.

## 5.2 PHARMACOKINETIC PROPERTIES

The pharmacokinetic profile of sotrovimab is consistent with a half-life extended IgG. The geometric mean  $C_{max}$  following a 1 hour IV infusion was 117.6 µg/mL (N = 290, CV% 46.5), and the geometric mean day 29 concentration was 24.5 µg/mL (N = 372, CV% 42.4) from all subjects with an available day 29 sample.

Based on population PK analyses, the PK of sotrovimab was not affected by age, sex, renal impairment, or mild or moderate hepatic impairment; body weight and BMI were significant covariates. Over the range from 40 to 160 kg, the effect of body weight on exposure was not considered clinically relevant and dose adjustment is not recommended.

### Absorption

No absorption studies have been conducted.

### Distribution

Based on noncompartmental analysis, the mean steady-state volume of distribution of sotrovimab was 8.1 L.

### Metabolism

Sotrovimab is an engineered human IgG1 monoclonal antibody degraded by proteolytic enzymes which are widely distributed in the body and not restricted to hepatic tissue.

### Excretion

Based on noncompartmental analysis, the mean systemic clearance (CL) was 125 mL/day, with a median terminal half-life of approximately 49 days.

## **Special patient populations**

### Children

The pharmacokinetics of sotrovimab in children under the age of 18 years have not been evaluated. However, the recommended dosing regimen in patients aged 12 years and older weighing at least 40 kg is expected to result in comparable serum exposures of sotrovimab as those observed in adults, based on an allometric scaling approach which accounted for effect of body weight changes associated with age on clearance and volume of distribution.

### Elderly

Based on population pharmacokinetic analysis, there was no difference in sotrovimab pharmacokinetics in elderly patients when compared with younger patients.

### Renal impairment

Sotrovimab, like other immunoglobulins, is too large to be excreted renally, thus renal impairment is not expected to have any effect on the elimination of sotrovimab. Furthermore, based on population pharmacokinetic analyses there was no difference in sotrovimab pharmacokinetics in patients with mild, moderate or severe renal impairment (eGFR <30 mL/min/1.73m<sup>2</sup>). There are limited data available in subjects with severe renal impairment.

### Hepatic impairment

Sotrovimab is degraded by widely distributed proteolytic enzymes, not restricted to hepatic tissue; therefore, changes in hepatic function are unlikely to have any effect on the elimination of sotrovimab. Furthermore, based on population pharmacokinetic analyses there is no difference in sotrovimab pharmacokinetics in patients with mild and moderate hepatic impairment. There are limited data available in patients with severe hepatic impairment.

## **5.3 PRECLINICAL SAFETY DATA**

### **Genotoxicity**

Genotoxicity studies have not been conducted with sotrovimab.

### **Carcinogenicity**

Carcinogenicity studies have not been conducted with sotrovimab.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 LIST OF EXCIPIENTS**

Histidine

Histidine hydrochloride monohydrate

Sucrose

Methionine

Polysorbate 80

Water for injections

## **6.2 INCOMPATIBILITIES**

XEVUDY concentrated injection solution for infusion must not be mixed with other medicinal products except those mentioned in section 4.2 DOSAGE AND METHOD OF ADMINISTRATION, Method of administration.

## **6.3 SHELF LIFE**

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG).

The expiry date can be found on the vial and packaging.

## **6.4 SPECIAL PRECAUTIONS FOR STORAGE**

### Unopened packs

Store refrigerated at 2°C to 8°C in the original carton. Do not freeze. Protect from light.

### Opened packs (in-use storage conditions)

The diluted solution of XEVUDY is intended to be used immediately. If immediate administration is not possible, the diluted solution may be stored at room temperature (up to 25°C) for up to 6 hours or refrigerated (2°C to 8°C) for up to 24 hours from the time of dilution until the end of administration (see Section 4.2 DOSAGE AND METHOD OF ADMINISTRATION, Method of Administration).

## **6.5 NATURE AND CONTENTS OF CONTAINER**

10 mL Type I clear glass vial, with a rubber stopper and flip-off aluminium over-seal.

XEVUDY is supplied as a single-use vial. Use in one patient on one occasion only. Contains no antimicrobial preservative.

## **6.6 SPECIAL PRECAUTIONS FOR DISPOSAL**

In Australia, any unused medicine or waste material should be disposed of by taking to your local pharmacy.

## **6.7 PHYSICOCHEMICAL PROPERTIES**

### **Chemical structure**

Not relevant

### **Cas number**

2423014-07-5

## **7 MEDICINE SCHEDULE (POISONS STANDARD)**

Schedule 4 – Prescription Only Medicine

## 8 SPONSOR

GlaxoSmithKline Australia Pty Ltd  
Level 4, 436 Johnston Street,  
Abbotsford, Victoria, 3067

## 9 DATE OF FIRST APPROVAL

20 August 2021

## 10 DATE OF REVISION

Not applicable

### SUMMARY TABLE OF CHANGES

Section Changed	Summary of new information

Version 1.0

Trademarks are owned by or licensed to the GSK.

© 2021 GSK group of companies or its licensor.