

# DALACIN<sup>®</sup> C Injection

Clindamycin phosphate

## Consumer Medicine Information

### What is in this leaflet

This leaflet answers some common questions about Dalacin C Injection. It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you being given Dalacin C Injection against the benefits this medicine is expected to have for you.

Dalacin C Injection should only be given to you by a health professional.

**If you have any concerns about using this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with your medicine.**

You may need to read it again.

### What DALACIN C Injection is used for

Dalacin C Injection is an antibiotic. It is used to treat infections in different parts of the body caused by bacteria.

This medicine works by killing or stopping the growth of bacteria causing your infection.

It will not work against viral infections such as colds or flu.

Dalacin C Injection is recommended for patients who are allergic to penicillin or patients for whom, in the judgment of the doctor, penicillin is inappropriate.

**Ask your doctor if you have any questions about why Dalacin C**

**Injection has been prescribed for you.**

Your doctor may have prescribed Dalacin C Injection for another reason.

This medicine is available only with a doctor's prescription.

Dalacin C Injection is not addictive.

### Before you are given DALACIN C Injection

**When you must not be given it**

**You must not be given Dalacin C Injection if you have an allergy to:**

- clindamycin or lincomycin
- any of the other ingredients listed at the end of this leaflet (see 'Product Description').

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

**You should not be given this medicine if the expiry date printed on the pack has passed or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal.

**If you are not sure whether you should be given this medicine, talk to your doctor.**

**Before you are given it**

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your doctor if you have or have had any of the following medical conditions:**

- severe diarrhoea associated with the use of antibiotics
- severe liver disease
- severe kidney disease
- bowel disease
- any gastrointestinal (stomach or gut) problems.

**Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.**

Your doctor can discuss with you the risks and benefits involved.

**If you have not told your doctor about any of the above, tell him/her before you are given Dalacin C Injection.**

### Children

Dalacin C Injection contains benzyl alcohol. Benzyl alcohol has been associated with a rare but serious side effect in infants. Your doctor will decide if treatment is appropriate.

### Taking other medicines

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and Dalacin C Injection may interfere with each other. These include:

- erythromycin, a medicine used to treat bacterial infections
- rifampicin, a medicine used to treat bacterial infections
- medicines used for muscle relaxation in anaesthesia.

These medicines may be affected by Dalacin C or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines. Your doctor will advise you.

**Tell your doctor or pharmacist if you are taking any of the following:**

- erythromycin, a medicine used to treat bacterial infections
- medicines used for muscle relaxation in anaesthesia.

Your doctor or pharmacist may have more information on medicines to be careful with or avoid while being given this medicine.

---

## How DALACIN C Injection is given

---

**Follow all directions given to you by your doctor and pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions in this leaflet, ask your doctor or pharmacist for help.**

### ***How much is given***

Your doctor will decide what dose of Dalacin C Injection you should receive and for how long you should receive it for. This depends on your condition, the infection being treated and how severe it is.

For children, the doctor will work out the dose based on their age, body weight and how severe the infection is.

### ***How it is given***

Dalacin C Injection is usually given as an injection into a vein

(intravenous drip) or into a muscle (intramuscular). It is given by a doctor or trained nurse.

Dalacin C Injection will be diluted if given as injection into a vein.

### ***If you take too much (overdose)***

As Dalacin C Injection is usually given to you in hospital under the close supervision of your doctor, it is unlikely that you will receive too much.

**However, if you experience severe side effects after being given Dalacin C Injection, tell your doctor or nurse immediately.**

Symptoms of an overdose are the same as those listed under the "Side effects" section but are usually of a more severe nature.

**Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital if you think that you or anyone else may have been given too much Dalacin C Injection.**

**Do this even if there are no signs of discomfort or poisoning.**

You may need urgent medical attention.

**Keep the telephone numbers for these services handy.**

**Have the Dalacin C Injection box or this leaflet available to give details if needed.**

---

## While you are being given DALACIN C Injection

---

### ***Things you must do***

**If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.**

**If you get severe diarrhoea, tell your doctor, pharmacist or nurse immediately. Do this even if it**

**occurs several weeks after Dalacin C Injection has been stopped.**

Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care.

**Do not take any medicines for diarrhoea without first checking with your doctor.**

**If you get a severe skin rash tell your doctor immediately. Do this even if the rash occurs after Dalacin C Injection has been stopped.**

**If you get a sore, white mouth or tongue while being given Dalacin C Injection or soon after Dalacin C Injection has been stopped, tell your doctor. Also tell your doctor if you get vaginal itching or discharge.**

This may mean you have a fungal/yeast infection called thrush. Sometimes the use of this medicine allows fungi/yeast to grow and the above symptoms to occur. Dalacin C Injection does not work against fungi/yeast.

**If you become pregnant while you are being given Dalacin C Injection, tell your doctor.**

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are being given Dalacin C Injection.**

**Tell all doctors, dentists and pharmacists who are treating you that you are being given Dalacin C Injection.**

### ***Things you must not do***

**Dalacin C Injection should not be given to treat any other complaints unless your doctor tells you.**

**Do not give your medicine to anyone else, even if they have the same condition as you.**

**Do not stop using your medicine without checking with your doctor.**

If you stop using Dalacin C Injection too soon, the infection may not clear

completely or your symptoms may return.

---

## Side effects

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are given Dalacin C Injection.**

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

It can be difficult to tell whether side effects are the result of taking Dalacin C Injection, effects of your condition or side effects of other medicines you may be taking. For this reason it is important to tell your doctor of any change in your condition.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Do not be alarmed by the list of side effects.**

You may not experience any of them.

**Tell your doctor or pharmacist if you notice any of the following and they worry you:**

- stomach pain, cramps or discomfort
- nausea and/or vomiting
- diarrhoea
- skin rash; irritation of the skin
- vaginal thrush - sore and itchy vagina and/or discharge
- low blood pressure (feeling of dizziness or light-headedness)
- joint pain and swelling
- pain, swelling, redness or formation of an abscess at the site of the injection
- loss or distorted sense of taste.

**Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you experience any of the following:**

- sudden signs of allergy such as rash, itching or hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing
- moderate or severe skin rash or blisters often with flu-like symptoms
- enlarged lymph glands and/or fever
- diarrhoea, usually with blood and mucus, stomach pain and fever
- yellowing of the eyes or skin, also called jaundice
- pain, swelling, redness and tenderness in vein or pain and swelling in leg
- chest pain, shortness of breath and/or fainting.

These are very serious side effects. You may need urgent medical attention or hospitalisation.

### **After finishing it**

**Tell your doctor immediately if you notice any of the following side effects, particularly if they occur several weeks after stopping treatment with Dalacin C Injection:**

- severe stomach cramps;
- watery and severe diarrhoea which may also be bloody;
- fever, in combination with one or both of the above.

Dalacin C Injection can cause some bacteria, which are normally present in the bowel and normally harmless to multiply and therefore cause the above symptoms. You may need urgent medical attention. However this side effect is rare.

Do not take any medicine for diarrhoea without first checking with your doctor.

**Tell your doctor if you notice anything else that is making you feel unwell.**

**Other side effects not listed above may also occur in some people.**

Some of these side effects (for example, abnormal blood test results and certain liver conditions) can only be found when your doctor does tests from time to time to check on your progress.

---

## After using DALACIN C Injection

---

### **Storage**

Dalacin C Injection will normally be stored in a hospital. It should be stored at 2°C to 8°C (Refrigerate. Do not freeze).

### **Disposal**

The hospital staff will dispose of any leftover Dalacin C Injection.

---

## Product description

---

### **What it looks like**

Dalacin C Injection appears as a clear colourless solution and comes in 2 mL or 4 mL glass ampoules.

### **Ingredients**

Dalacin C Injection contains clindamycin 2-phosphate as the active ingredient

It also contains:

- benzyl alcohol
- disodium edetate
- sodium hydroxide
- hydrochloric acid
- water for injections.

### **Supplier**

Dalacin C Injection is supplied in Australia by:

Pfizer Australia Pty Ltd  
Sydney NSW 2000

Toll Free number: 1800 675 229

www.pfizer.com.au

® = Registered Trademark

© Copyright

This leaflet was prepared in February  
2021.

***Australian registration  
numbers***

300 mg/2 mL injection: AUST R  
12294

600 mg/4 mL injection: AUST R  
47636