

# **AUSTRALIAN PRODUCT INFORMATION – MERSYNOFEN (IBUPROFEN AND PARACETAMOL) FILM COATED TABLETS**

## **1 NAME OF THE MEDICINE**

Ibuprofen 200mg and paracetamol 500mg film coated tablets

## **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Ibuprofen 200mg and paracetamol 500mg

## **3 PHARMACEUTICAL FORM**

Film coated tablet

## **4 CLINICAL PARTICULARS**

### **4.1 THERAPEUTIC INDICATIONS**

Mersynofen is indicated for the temporary relief of acute (short term) pain and/or inflammation associated with headache, migraine headache, tension headache, sinus pain, toothache, dental procedures, backache, muscular aches and pains, period pain, sore throat, tennis elbow, rheumatic pain and arthritis, and the aches and pains associated with colds and flu.

### **4.2 DOSE AND METHOD OF ADMINISTRATION**

Adults under 65 and children from 12 years: 1 tablet every 8 hours as necessary (maximum 3 tablets in 24 hours). Do not split the tablets. Keep to the recommended dose.

Mersynofen should not be used for more than 3 days at a time (or not more than 2 days at a time for adolescents aged 12 to 17 years).

- Not recommended for children under 12 years of age.
- Not recommended for adults 65 years and over.

Monitoring advice: if symptoms persist please consult your healthcare professional.

The lowest effective dose should be used for the shortest duration necessary to relieve symptoms (see under section 4.4 Special warnings and precautions for use).

### **4.3 CONTRAINDICATIONS**

This product is contraindicated:

- In patients with a known hypersensitivity to ibuprofen, paracetamol or any other constituent of the medicinal product

- In patients with a history of hypersensitivity reactions (e.g. bronchospasm, angioedema, rhinitis or urticaria) associated with aspirin or other anti-inflammatory drugs or analgesic drugs.
- In patients with a history of, or an existing gastrointestinal ulceration/perforation or bleed, or other stomach disorder.
- In patients with impaired hepatic function, impaired renal function or heart failure
- In patients with asthma
- In pregnancy
- In patients with conditions that predispose to renal failure
- In concomitant use with ibuprofen or other NSAID-containing products, including cyclooxygenase-2 (COX-2) specific inhibitors and aspirin or other anti-inflammatories as there is an increased risk of adverse reactions
- In concomitant use with other paracetamol-containing products as there is an increased risk of serious adverse effects; patients should be advised not take with any other paracetamol containing products. Immediate medical advice should be sought if this occurs, even if they feel well, as this can result in an overdose.
- In patients aged 65 years and over and in children under 12 years
- In patients with severe dehydration (caused by vomiting, diarrhoea or insufficient fluid intake)

#### **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE**

##### **Identified precautions**

The hazard of paracetamol overdose is greater in patients with non-cirrhotic alcoholic liver disease. Immediate medical advice should be sought in the event of an overdose, even if the patient feels well, because of the risk of delayed, serious liver damage. Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms.

##### ***Diabetes***

Caution is required in patients suffering from diabetes. Paracetamol falsely elevates continuous blood glucose monitor (CGM) readings compared to finger stick (BG meter) readings. This is applicable for those using CGM devices with or without an automated insulin delivery pump e.g. in type I diabetes.

##### ***Respiratory disorders***

Caution is required in patients with a history of bronchial asthma or allergic disease since NSAIDs have been reported to precipitate bronchospasm. The product is contraindicated in asthma (see under section 4.3 Contraindications above).

##### ***Cardiovascular and cerebrovascular effects***

Observational studies have indicated that NSAIDs may be associated with an increased risk of serious cardiovascular events, including myocardial infarction and stroke, which may increase with dose or duration of use.

Patients with cardiovascular disease, history of atherosclerotic cardiovascular disease or cardiovascular risk factors may also be at greater risk.

Patients should be advised to remain alert for such cardiovascular events, even in the absence of previous cardiovascular symptoms. Patients should be informed about signs and/or symptoms of serious cardiovascular toxicity and the steps to take if they occur.

Appropriate monitoring and advice are required for patients with a history of hypertension as fluid retention and oedema have been reported in association with NSAID therapy. The product is contraindicated in patients with heart failure (see under section 4.3 Contraindications above).

Clinical trial data suggest that the use of ibuprofen, particularly at high doses (2400 mg daily) may be associated with an increased risk of arterial thrombotic events (e.g. myocardial infarction or stroke). Overall, epidemiological studies do not suggest that low dose ibuprofen (e.g. <1200mg daily) is associated with an increased risk of myocardial infarction.

Patients with uncontrolled hypertension, established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease should only be treated with FDC after careful consideration. Similar consideration should be made before initiating treatment for patients with risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes mellitus and smoking). The product is contraindicated in heart failure (see under section 4.3 Contraindications above).

### ***Gastrointestinal bleeding, ulceration and perforation***

Gastrointestinal (GI) bleeding, ulceration and perforation, which can be fatal, has been reported with all NSAIDs at any time during treatment, with or without warning symptoms or a previous history of serious GI events.

Caution should be advised in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as oral corticosteroids, anticoagulants such as warfarin, selective serotonin-reuptake inhibitors (SSRIs), or antiplatelet agents.

The product is contraindicated in patients with a history of GI toxicity including ulceration (see under section 4.3 Contraindications above).

When GI bleeding or ulceration occurs in patients receiving FDC, the treatment should be withdrawn.

### ***SLE and mixed connective tissue disease***

In patient with systemic lupus erythematosus (SLE) and mixed connective tissue disease disorders there may be an increased risk of aseptic meningitis.

### ***Dermatological***

Serious skin reactions, some of them fatal, including bullous and exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis (TEN), have been reported very rarely in association with the use of NSAIDs and paracetamol. Patients appear to be at highest risk of these reactions early in the course of therapy, the onset of the reaction occurring in the majority of cases within the first month of treatment. Use of this product should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

### ***Masking of symptoms of underlying infections***

Mersynofen can mask symptoms of infection, which may lead to delayed initiation of appropriate treatment and thereby worsening the outcome of the infection. This has been observed in bacterial community acquired pneumonia and bacterial complications to varicella. When Mersynofen is administered for fever or pain relief in relation to infection, monitoring of infection is advised. In non-hospital settings, the patient should consult a doctor if symptoms persist or worsen.

### **Use in renal and hepatic impairment**

The administration of NSAIDs may cause a dose-dependent reduction in prostaglandin formation and precipitate renal failure. Patients at greatest risk of this reaction are those with impaired renal function, cardiac impairment, liver dysfunction, those taking diuretics and the elderly. The product is contraindicated in patients with impaired renal or liver function or heart failure and in patients 65 years of age or older (see under section 4.3 Contraindications above). Renal function should be monitored in other at risk patients.

As with other NSAIDs, elevations of one or more liver function tests may occur in up to 15% of patients. These abnormalities may progress, may remain essentially unchanged or may resolve with continued therapy. Meaningful elevations (three times the upper limit of normal) of ALT or AST occurred in controlled clinical trials in less than 1% of patients.

Patients should be advised to remain alert for hepatotoxicity and be informed about the signs and/or symptoms of hepatotoxicity (e.g. nausea, fatigue, lethargy, pruritus, jaundice, abdominal tenderness in the right upper quadrant and “flu-like” symptoms).

### **Use in the elderly**

Mersynofen is contraindicated in adults aged 65 years and over.

### **Paediatric use**

Mersynofen is contraindicated in children under 12 years of age since no investigations have been carried out with this product in this age group.

There is a risk of renal impairment in dehydrated adolescents.

### **Effects on laboratory tests**

No information is available regarding Mersynofen and laboratory tests.

## **4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS**

This product is contraindicated in combination with:

- Aspirin
- Other paracetamol containing products
- Other NSAIDs including cyclo-oxygenase-2 selective inhibitors
- Other anti-inflammatories and analgesics as concomitant use may increase the risk of adverse reactions.

This product (like any other paracetamol containing products) should be used with caution in combination with:

- Chloramphenicol: increased plasma concentration of chloramphenicol
- Cholestyramine: the speed of absorption of paracetamol is reduced by cholestyramine
- Therefore, cholestyramine should not be taken within one hour if maximal analgesia is required
- Metoclopramide and domperidone: the absorption of paracetamol is increased by metoclopramide and domperidone. However, concurrent use need not be avoided
- Warfarin: the anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular use of paracetamol with increased risk of bleeding; occasional doses have no significant effect

This product (like any other ibuprofen containing products and NSAIDs) should be used with caution in combination with:

- Anticoagulants: NSAIDs may enhance the effects of anticoagulants, i.e. warfarin
- Antihypertensives: NSAIDs may reduce the effects of these drugs
- Antiplatelet agents and selective serotonin reuptake inhibitors (SSRIs): increased risk of gastrointestinal bleeding
- Cardiac glycosides: NSAIDs may exacerbate cardiac failure, reduce glomerular filtration rate (GFR) and increase plasma glycoside levels
- Ciclosporin: increased risk of nephrotoxicity
- Corticosteroids: Increased risk of gastrointestinal ulceration or bleeding
- Diuretics: reduced diuretic effect. Diuretics may increase the risk of nephrotoxicity of NSAIDs
- Lithium: decreased elimination of lithium
- Methotrexate: decreased elimination of methotrexate
- Mifepristone: NSAIDs should not be used for 8-12 days after mifepristone administration as NSAIDs can reduce the effect of mifepristone
- Quinolone antibiotics: animal data indicate that NSAIDs can increase the risk of convulsions associated with quinolone antibiotics. Patients taking NSAIDs and quinolones may have an increased risk of developing convulsions
- Tacrolimus: possible increased risk of nephrotoxicity when NSAIDs are given with tacrolimus
- Zidovudine: increased risk of haematological toxicity when NSAIDs are given concomitantly with zidovudine. There is evidence of an increased risk of hemarthrosis and haematoma in HIV+ haemophiliacs receiving concurrent treatment with zidovudine and ibuprofen.

#### **4.6 FERTILITY, PREGNANCY AND LACTATION**

##### **Effects on fertility**

The use of the product may impair female fertility and is not recommended in women attempting to conceive.

##### **Use in pregnancy**

Category C

Drugs which owing to their pharmacological effects have caused or may be suspected of causing harmful effects on the human foetus or neonate without causing malformation. These effects may be reversible.

There is no experience of use of this product in humans during pregnancy. Therefore this product is contraindicated for use during pregnancy.

Congenital abnormalities have been reported in association with NSAID administration in man; however these are low in frequency and do not appear to follow any discernible pattern. Use of NSAIDs during the last trimester of pregnancy may cause effects on the foetal cardiovascular system (risk of closure of ductus arteriosus), and the onset of labour may be delayed and the duration increased with an increased bleeding tendency in both mother and child.

Epidemiological studies in human pregnancy have shown no ill effects due to paracetamol use at the recommended dosage.

## **Use in lactation**

Ibuprofen and its metabolites can pass in very small amounts (0.0008% of the maternal dose) into the breast milk. No harmful effects to infants are known.

Paracetamol is excreted in breast milk but not in a clinically significant amount. Available published data do not contraindicate breastfeeding.

Therefore it is not necessary to interrupt breastfeeding for short term treatment with the recommended dose of this product.

## **4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES**

No data available.

## **4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)**

Clinical trials with this product have not indicated any other undesirable effects other than those for ibuprofen or paracetamol alone.

In short term clinical trials, FDC was shown to have a safety profile comparable to that of placebo. The body systems most commonly affected by ADE associated with FDC treatment were the gastrointestinal and nervous systems. The most commonly reported ADEs were nausea, vomiting, diarrhoea, dyspepsia, dizziness and headache.

One study recorded ADEs and abnormal haematological and biochemical results in subjects treated with FDC three times daily for 13 weeks. The median incidence of all ADE considered to be moderate and severe, regardless of causality, was 1.1 per person-days exposure in all four treatment groups. The commonest treatment related ADEs were dyspepsia, diarrhoea and nausea.

Compared to the other treatments, more incidence of diarrhoea were reported in association with FDC; a higher incidence of liver function abnormalities with paracetamol; and a higher incidence of early but transient abnormal liver tests with paracetamol vs ibuprofen. At the end of the study (13 weeks) the incidence of treatment related ADEs was significantly higher in patients taking one (50.5%) or 2 (51.3%) FDC compared to ibuprofen (42%,  $p = 0.04$ ), but not paracetamol (45.5%).

Mean haemoglobin decreased in all groups throughout the study. At study end, the proportion of patients experiencing a decrease of 2g/100mL or greater decrease in haemoglobin was significantly higher in the 2 FDC group (6.9%) compared to those taking paracetamol (0.9%,  $p = 0.011$ ), ibuprofen (0.9%,  $p = 0.001$ ) and one FDC (1.8%,  $p = 0.0096$ ).

For FDC in general the percentage of subjects who experience an adverse event, as well as the range of ADE observed, are similar to those established for ibuprofen and paracetamol when administer alone.

In clinical trials, the product is administered in single or multiple doses.

The following is a list of adverse effects from pharmacovigilance data experienced by patients taking ibuprofen alone or paracetamol alone in short term and long term use.

Adverse events may be minimized by using the minimum effective dose for the shortest duration necessary to control symptoms.

### **Common (occurring >1% and <10%)**

Gastrointestinal: abdominal pain, diarrhoea, dyspepsia, nausea, stomach discomfort, flatulence and constipation and vomiting. Investigations: alanine aminotransferase increased, gamma

glutamyl transferase increased and liver function tests abnormal with paracetamol. Blood creatinine increased and blood urea increased.

**Uncommon (occurring >0.1% and <1%)**

Gastrointestinal: peptic ulcer, perforation or gastrointestinal haemorrhage, with symptoms of melaena, haematemesis sometimes fatal, particularly in the elderly. Ulcerative stomatitis and exacerbation of ulcerative colitis and Crohn's disease. Less frequently gastritis has been observed and pancreatitis reported.

Skin and subcutaneous tissue disorders: rashes of various types (including urticarial) and pruritis. Angioedema and swelling face.

Investigations: aspartate aminotransferase increased, blood alkaline phosphatase increased, blood creatine phosphokinase increased, blood creatinine increased, haemoglobin decreased and platelet count increased.

Nervous system disorders: headache and dizziness.

Eye disorders: visual disturbance.

**Very rare (occurring < 0.01%)**

Blood and lymphatic system disorders: haematopoietic disorders (agranulocytosis, anaemia, aplastic anaemia, haemolytic anaemia, leucopaenia, neutropaenia, thrombocytopaenia and pancytopaenia). First signs are fever, sore throat, superficial mouth ulcers, flu-like symptoms, severe exhaustion, unexplained bleeding and bruising and nose bleeds.

Immune system disorders: hypersensitivity reactions have been reported. These may consist of non-specific allergic reactions and anaphylaxis. Symptoms of severe hypersensitivity reactions can include facial, tongue and larynx swelling, dyspnoea, tachycardia, hypotension, anaphylaxis, angioedema or severe shock.

Psychiatric disorders: confusion, depression and hallucinations.

Nervous system disorders: paraesthesia, optic neuritis and somnolence. Single cases of aseptic meningitis in patients with existing autoimmune disorders (e.g. systemic lupus erythematosus and mixed connective tissue disease) during treatment with ibuprofen, with symptoms such as stiff neck, headache, nausea, vomiting, fever or disorientation have been observed.

Ear and labyrinth disorders: tinnitus and vertigo.

Cardiac disorders: hypertension and cardiac failure have been reported in association with NSAID treatment. Clinical trial and epidemiological data suggest that use of ibuprofen, particularly at high doses (2400 mg daily), and in long term treatment may be associated with a small increased risk of arterial thrombotic events (e.g. myocardial infarction or stroke).

Respiratory, thoracic and mediastinal disorders: respiratory reactivity including asthma, exacerbation of asthma, bronchospasm and dyspnoea.

Hepatobiliary disorders: abnormal liver function, hepatitis and jaundice. In overdose, paracetamol can cause acute hepatic failure, hepatic failure, hepatic necrosis and liver injury.

Skin and subcutaneous tissue disorders: hyperhidrosis, purpura and photosensitivity. Exfoliative dermatoses. Bullous reactions including bullous erythema multiforme, Stevens-Johnson syndrome and toxic epidermal necrolysis.

Renal and urinary disorders: oedema, nephrotoxicity in various forms, including interstitial nephritis, nephrotic syndrome, and acute and chronic renal failure.

General disorders and administration site conditions: fatigue and malaise.

Hypersensitivity reactions have been reported following treatment with both paracetamol and ibuprofen. These may consist of:

- a) Non-specific allergic reactions and anaphylaxis.
- b) Respiratory tract reactivity e.g. asthma, aggravated asthma, bronchospasm and dyspnoea.
- c) Assorted skin disorders, including rashes of various types, pruritus, urticaria, purpura, angioedema and more rarely bullous dermatoses (including toxic epidermal necrolysis and bullous erythema multiforme).

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems) (Australia).

#### **4.9 OVERDOSE**

For general advice on overdose management, contact the Poisons Information Centre, telephone number 13 11 26 (Australia).

##### **Paracetamol**

Liver damage is possible in adults who have taken 10 g (equivalent to 20 tablets) or more of paracetamol. Ingestion of 5 g (equivalent to 10 tablets) or more of paracetamol may lead to liver damage if the patient has one or more of the risk factors below:

- a) Is on long term treatment with carbamazepine, phenobarbitone, phenytoin, primidone, rifampicin, St John's Wort or other drugs that induce liver enzymes.
- b) Regularly consumes alcohol in excess of recommended amounts.
- c) Is likely to be glutathione depleted e.g. eating disorders, cystic fibrosis, HIV infection, starvation, cachexia.

##### ***Symptoms***

Symptoms of paracetamol overdose in the first 24 hours include pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion as liver function tests become abnormal. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning, hepatic failure may progress to encephalopathy, disseminated intravascular coagulation, haemorrhage, hypoglycaemia, cerebral oedema and death. Acute renal failure with acute tubular necrosis, strongly suggested by loin pain, haematuria and proteinuria, may develop even in the absence of severe liver damage. Cardiac arrhythmias and pancreatitis have been reported.

##### ***Management***

Immediate treatment is essential in the management of paracetamol overdose. Despite a lack of significant early symptoms, patients should be referred to hospital urgently for immediate medical attention. Symptoms may be limited to nausea or vomiting and may not reflect the severity of overdose or the risk of organ damage. Management should be in accordance with established treatment guidelines.

Treatment with activated charcoal should be considered if the overdose has been taken within 1 hour. Plasma paracetamol concentration should be measured at 4 hours or later after ingestion (earlier concentrations are unreliable).

Treatment with N-acetylcysteine may be used up to 24 hours after ingestion of paracetamol, however the maximum protective effect is obtained up to 8 hours post- ingestion. The effectiveness of the antidote declines sharply after this time.



If required, the patient should be given intravenous N-acetylcysteine in line with the established dosage schedule. If vomiting is not a problem, oral methionine may be a suitable alternative for remote areas, outside hospital.

Patients who present with serious hepatic dysfunction beyond 24 hours from ingestion should be managed in accordance with established guidelines.

## **Ibuprofen**

### ***Symptoms***

Most patients who have ingested clinically important amounts of NSAIDs will develop no more than nausea, vomiting, epigastric pain, or more rarely diarrhoea. Tinnitus, headache and gastrointestinal bleeding are also possible. In more serious poisoning, toxicity is seen in the central nervous system, manifesting as drowsiness, occasionally excitation and disorientation or coma. Occasionally patients develop convulsions. In serious poisoning, metabolic acidosis may occur and prolong the prothrombin time (PT) and increase the international normalised ratio (INR), probably due to interference with the actions of circulating clotting factors. Acute renal failure and liver damage may occur if there is co-incident dehydration. Exacerbation of asthma is possible in asthmatics.

### ***Management***

Management should be symptomatic and supportive and include the maintenance of a clear airway and monitoring of cardiac and vital signs until stable. Consider oral administration of activated charcoal if the patient presents within 1 hour of ingestion of a potentially toxic amount. If frequent or prolonged, convulsions should be treated with intravenous diazepam or lorazepam. Give bronchodilators for asthma.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 PHARMACODYNAMIC PROPERTIES**

#### ***Mechanism of action***

The pharmacological actions of ibuprofen and paracetamol differ in their site and mode of action. These complementary modes of action result in greater antinociception than the single actives alone.

Ibuprofen possesses analgesic, antipyretic and anti-inflammatory properties, similar to other non-steroidal anti-inflammatory drugs (NSAIDs). Its mechanism of action is unknown, but it is thought to be through peripheral inhibition of cyclooxygenases and subsequent prostaglandin synthetase inhibition.

Paracetamol is a para-aminophenol derivative that exhibits analgesic and antipyretic activity.

Paracetamol has minimal anti-inflammatory action. The precise mechanism of action remains uncertain; it is believed to include inhibition of prostaglandin synthesis, primarily within the central nervous system.

#### ***Clinical trials***

#### **Preclinical safety data**

The toxicological safety profiles of ibuprofen and paracetamol individually have been established in animal experiments and in humans from extensive clinical experience.

## Summary of clinical data

Five randomised, double-blind, placebo-controlled studies were conducted to evaluate the efficacy and safety of 200mg/500mg ibuprofen/500 mg paracetamol fixed dose combination (FDC) when used to treat post-operative dental pain, pain associated with dysmenorrhoea and chronic knee pain.

### **Study 1**

This efficacy study was a two part study (a single dose phase and a multiple dose phase). Seven hundred and thirty five subjects with post-operative dental pain were randomised to one of eight treatment groups in Part 1 (placebo, ibuprofen 200 mg or 400 mg, paracetamol 500 mg or 1000 mg, ibuprofen 100 mg plus paracetamol 250 mg (½ tablet FDC), ibuprofen 200 mg plus paracetamol 500 mg (1 tablet FDC) or 400 mg ibuprofen plus 1000 mg paracetamol (2 tablets FDC).

In Part 1 of the study (single-dose phase), the primary efficacy variable was the mean differences in the sum of total pain relief and pain intensity difference (SPRID 0-8h) for pairwise comparisons.

Primary efficacy variable comparisons during Part 1 (single dose) all favoured the combination product over the comparators – that is, 2 tablets of FDC were more effective than 400 mg ibuprofen, 1000 mg paracetamol or placebo, and 1 tablet of FDC was more effective than 200 mg ibuprofen, 500 mg paracetamol or placebo. The majority of the secondary efficacy endpoints (including pain relief intensity difference 8 hours post dose, subjects overall assessment of medication, time to meaningful pain relief, duration of effects, and total pain relief over 8 hours) were consistent with the primary efficacy findings.

Seven hundred and fifteen subjects entered Part 2 (multiple dose phase) of the pivotal study, which involved only combinations – ½, 1 or 2 tablets of FDC – (no single actives) against placebo. The primary efficacy endpoint was the number of completed 24-hour periods with no more than one dose of rescue medication and with the subject's overall assessment always rated as at least good, in subjects who had taken the combination treatment or placebo in both parts 1 and 2 of the study.

One or two tablets of FDC were statistically significantly superior to placebo for the primary efficacy endpoint. The secondary efficacy variables (including time to treatment failure, duration between doses, peak pain relief and median score for subjects overall assessment) showed mixed results, with 1 tablet of FDC not significantly different to placebo for all parameters.

In Part 1, subjects taking either the 1 or 2 tablet doses of FDC experienced significantly fewer adverse effects than the placebo group, and subjects taking the 1 tablet dose of FDC also experienced significantly fewer adverse events than the 500 mg paracetamol group, with no significant differences in adverse events between any other groups. For the study overall, there were no significant differences in adverse events between any treatment groups. The most common adverse events in all groups were swelling face, nausea, vomiting and headache.

### **Study 2**

An exploratory, single dose, efficacy and safety study in 234 subjects with post-operative dental pain was also conducted. The double blind, double-dummy study compared 400 mg ibuprofen plus 1000 mg paracetamol (equivalent to 2 tablets FDC) with 200 mg ibuprofen plus 500 mg paracetamol (equivalent to 1 tablet FDC), 400 mg ibuprofen, 1000 mg paracetamol and placebo. Both doses of the combination treatment were significantly more efficacious as assessed by the primary efficacy parameter, SPRID (0-8 hr) than placebo and 1000 mg paracetamol. The higher

dose combination (equivalent to 2 tablets of FDC), but not the lower dose combination (equivalent to 1 tablet FDC), was significantly more efficacious than 400 mg ibuprofen.

Both combination treatments were significantly more efficacious than placebo for the majority of secondary efficacy variables (including total pain relief (TOTPAR), sum of pain intensity difference (SPID) and SPRID over 0-4, 0-6 and 0-8 hours, peak pain relief and time to pain relief). The secondary efficacy variables showed mixed results for the comparisons of the combination treatments with ibuprofen 400 mg and paracetamol 1000 mg.

Each of the treatments was well tolerated and the adverse event profiles of the combination treatments were comparable to that of either drug administered alone.

### **Study 3**

A double-blind, single dose, placebo-controlled, randomised study compared 1 or 2 tablets of FDC with a combination of paracetamol 1000 mg plus codeine 30 mg (2 tablets Panadeine Extra®) and ibuprofen 400 mg plus codeine 25.6 mg (2 tablets Nurofen Plus®) in 678 subjects with post-operative dental pain.

The study was conducted in subjects >16 years of age with moderate to severe pain.

The primary efficacy endpoint was the sum of the mean scores of pain relief (PR) combined with pain intensity (PI) differences over 12 hours (SPRID 0–12 h), i.e. the sum of the PI difference and the PR score integrated over the follow-up time period).

This study showed that for the primary efficacy variable, after a single dose over a 12 hour evaluation period, 1 tablet of FDC was statistically significantly more efficacious than 2 tablets of Panadeine Extra® or placebo, and non-inferior in analgesic effect to 2 tablets of Nurofen Plus®. Similar results were observed for the majority of the secondary efficacy parameters (which included SPRID over 4, 6 and 8 hours, SPID over 4, 6, 8 and 12 hours, TOTPAR over 4, 6, 8 and 12 hours, peak pain relief and pain intensity difference, subjects overall assessment, and time of onset and duration of action), although there were no differences between any of the active treatment groups in time to meaningful pain relief.

Fewer treatment emergent, and treatment-related treatment emergent adverse events occurred with both 2 tablets of FDC (ibuprofen 400 mg plus paracetamol 1000 mg) and 1 tablet of FDC (ibuprofen 200 mg plus paracetamol 500 mg) compared with Nurofen Plus®, Panadeine Extra® and placebo. All these comparisons achieved statistical significance with the exception of 1 tablet of FDC when compared with placebo. The adverse event profile was consistent with patients having undergone third molar extraction and no safety issues were raised.

An additional study examined the efficacy and safety of the combination on an alternative pain model, primary dysmenorrhoea.

### **Study 4**

A double blind, randomised, crossover, single dose, single centre study examined the analgesic efficacy and tolerability of FDC in 94 subjects with primary dysmenorrhoea. Subjects received one of the following treatments: FDC (1 tablet) + placebo (1 tablet); FDC (2 tablets); or placebo (2 tablets). The study was conducted in females >18 years of age with primary dysmenorrhoea with moderate to severe cramping pain in at least 4 of the previous 6 months.

The primary efficacy endpoint was total pain relief over 0-6 hours (TOTPAR<sub>6</sub>). Secondary endpoints included TOTPAR over 2 and 4 hours, SPRID over 2, 4 and 6 hours, SPID over 2, 4 and 6 hours, and subject's overall assessment of medication.

Two tablets of FDC were statistically more efficacious (TOTPAR<sub>0-6h</sub>) than placebo (p=0.0001), and approached significance for one FDC compared to placebo (p=0.054). Two

tablets of FDC provided significantly greater pain relief compared to placebo from 2 hours post dose onward ( $p \leq 0.01$  all-time points), and one tablet FDC provided significantly more pain relief than placebo at the 4 ( $p < 0.05$ ) and 6 hour ( $p < 0.01$ ) assessment point.

The percentage of patients who rated their study medicine as ‘good’, ‘very good’ or ‘excellent’ was 63.3% for 2 FDC, 57.1% for one FDC and 43.3% for placebo. The corresponding percentage of patients rating their medicines as ‘poor’ was respectively 12.2%, 15.3% and 31.1%.

There were no withdrawals due to adverse events. Both the higher and lower dose combinations were well tolerated. The incidence of events did not differ with either treatment compared to placebo. Eleven patients reported 14 events (13 mild, 1 moderate) after taking the lower dose combination, seven patients reported 7 events (all mild) after taking the higher dose combination and nine patients reported 13 events (7 mild, 6 moderate) after taking placebo. There were no clinically significant laboratory abnormalities and no changes in vital signs during the course of the study.

## Study 5

This study evaluated a total of 892 patients (mean age 60.6 years, 49% female) experiencing knee pain, but who were not under medical supervision for this condition. Subjects had to be aged 40 years or older, experienced knee pain for most of the past 3 months and on 4 of the 7 preceding days. They were randomized to receive one of the following four treatments: 2 capsules of a combination (FDC) containing ibuprofen 200mg/paracetamol 500mg; one FDC capsule and one placebo capsule; 2 capsules of ibuprofen 200mg; or 2 capsules of paracetamol 500mg.

The primary short term efficacy endpoint was the difference between treatment groups in the Western Ontario McMaster Universities osteoarthritis index (WOMAC) pain scale (normalized to 0 to 100mm scale) after 10 day’s treatment. The primary long term efficacy endpoint was the patient global assessment (PG) of study medicine after 13 week’s treatment. The secondary endpoints analysed were the change in physical function from baseline, change in stiffness from baseline, change in composite score from baseline, time taken (seconds) from sitting to standing and acceptability of knee pain in the last 48 hours.

Two tablets of FDC were statistically more efficacious than 1000mg of paracetamol after 10 days of treatment ( $p = 0.0012$ ). This difference was maintained to 13 weeks ( $p < 0.001$ ). One FDC showed a non-significant benefit in pain relief over paracetamol. After 13 weeks of treatment patient global assessment rated one or two FDC as at least “excellent” or “good”, when compared to 1000mg paracetamol ( $p = 0.0152$  and  $p = 0.0002$  respectively). One FDC was associated with significantly better scores than paracetamol in physical functioning ( $p = 0.04$ ) and role-physical score ( $p = 0.0014$ ) at week 7 and in social functioning ( $p = 0.03$ ) at week 13. There were no significant differences between 400 mg ibuprofen and either 1 or 2 FDC for any efficacy parameter.

In this study, the median incidence of all ADE considered to be moderate and severe, regardless of causality, was 1.1 per person-days exposure in all four treatment groups. The commonest treatment related ADEs were dyspepsia, diarrhoea and nausea.

Compared to the other treatments, more incidence of diarrhoea were reported in association with FDC; a higher incidence of liver function abnormalities with paracetamol; and a higher incidence of early but transient abnormal liver tests with paracetamol vs ibuprofen. At the end of the study (13 weeks) the incidence of treatment related ADEs was significantly higher in patients taking one (50.5%) or 2 (51.3%) FDC compared to ibuprofen (42%,  $p = 0.04$ ), but not paracetamol (45.5%).

Mean haemoglobin decreased in all groups throughout the study. At study end, the proportion of patients experiencing a decrease of 2g/100mL or greater decrease in haemoglobin was significantly higher in the 2 FDC group (6.9%) compared to those taking paracetamol (0.9%,  $p = 0.011$ ), ibuprofen (0.9%,  $p = 0.001$ ) and one FDC (1.8%,  $p = 0.0096$ ).

## **5.2 PHARMACOKINETIC PROPERTIES**

### ***Absorption***

Ibuprofen is well absorbed from the gastrointestinal tract and is extensively bound to plasma proteins. Ibuprofen diffuses into the synovial fluid. Plasma levels of ibuprofen from this product are detected from 5 minutes with peak plasma concentrations achieved within 1-2 hours after ingestion on an empty stomach. When taken with food, peak plasma levels are delayed by a median of 25 minutes, but the overall extent of absorption is equivalent.

Paracetamol is readily absorbed from the gastrointestinal tract. Plasma protein binding is negligible at usual therapeutic concentrations, although this is dose-dependent. Plasma levels of paracetamol from this product are detected from 5 minutes with peak plasma concentrations occurring at 0.5-0.67 hours after ingestion on an empty stomach.

### ***Metabolism***

Ibuprofen is metabolised in the liver to two major metabolites with primary excretion via the kidneys, either as such or as major conjugates, together with a negligible amount of unchanged ibuprofen.

Paracetamol is metabolised in the liver and excreted in the urine mainly as the glucuronide and sulphate conjugates, with about 10% as glutathione conjugates. A minor hydroxylated metabolite, which is usually produced in very small amounts by mixed function oxidases in the liver and detoxified by conjugation with liver glutathione, may accumulate following paracetamol overdose and cause liver damage.

### ***Excretion***

Excretion of ibuprofen by the kidney is both rapid and complete. The elimination half-life is approximately 2 hours.

Less than 5% of paracetamol is excreted as unchanged paracetamol. The elimination half-life is approximately 3 hours.

No significant differences in the paracetamol or ibuprofen pharmacokinetic profiles are observed in the elderly.

The bioavailability and pharmacokinetic profiles of ibuprofen and paracetamol taken individually are not altered when taken in combination as a single or repeat dose.

## **5.3 PRECLINICAL SAFETY DATA**

### ***Genotoxicity***

No information is available regarding Mersynofen and genotoxicity.

### ***Carcinogenicity***

No information is available regarding Mersynofen and carcinogenicity.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 LIST OF EXCIPIENTS

maize starch  
magnesium stearate  
microcrystalline cellulose  
pregelatinised maize starch  
croscarmellose sodium  
sodium lauryl sulfate  
purified talc  
opadry QX Quick and Flexible Film Coating System 321A180025 White  
opadry fx special effects film coating system 63F97546 Silver

### 6.2 INCOMPATIBILITIES

Incompatibilities were either not assessed or not identified as part of the registration of this medicine.

### 6.3 SHELF LIFE

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

### 6.4 SPECIAL PRECAUTIONS FOR STORAGE

Store below 30°C.

### 6.5 NATURE AND CONTENTS OF CONTAINER

Container type:

Mersynofen tablets are packed in PVC/PVDC/Al blister packs.

Pack sizes of Mersynofen tablets: 2, 3, 6, 10, 12, 16, 24 or 30 tablets\*



\*Not all pack sizes will be marketed

### 6.6 SPECIAL PRECAUTIONS FOR DISPOSAL

In Australia, any unused medicine or waste material should be disposed of by taking to your local pharmacy.

### 6.7 PHYSICOCHEMICAL PROPERTIES

#### *Chemical structure*



Ibuprofen

Molecular formula: C<sub>13</sub>H<sub>18</sub>O<sub>2</sub>

MW: 206.3

**CAS number**

15687-27-1

Paracetamol

Molecular formula: C<sub>8</sub>H<sub>9</sub>NO<sub>2</sub>

MW: 151.16

103-90-2

## **7 MEDICINE SCHEDULE (POISONS STANDARD)**

2, 3, 6, 10, 12 and 16 tablets: Pharmacy Medicine (S2)

24 and 30 tablets: Pharmacist Only Medicine (S3)

## **8 SPONSOR**

Sanofi Consumer Healthcare,  
87 Yarraman Place  
Virginia, QLD 4014,  
Australia

## **9 DATE OF FIRST APPROVAL**

09 August 2019

## **10 DATE OF REVISION**

16 October 2020

### **SUMMARY TABLE OF CHANGES**

<b>Section Changed</b>	<b>Summary of new information</b>
<b>4.2</b>	Addition of information on using the lowest effective dose should be used for the shortest duration
<b>4.3, 4.4, 4.8, 4.9</b>	Addition of safety information including a warning in relation to masking of symptoms