Minims[®] Oxybuprocaine Eye Drops

Oxybuprocaine hydrochloride Eye Drops

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Minims Oxybuprocaine.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you using Minims
Oxybuprocaine against the benefits they expect it will have for you.

If you have any concerns about this medicine, ask your doctor.

Keep this leaflet in a safe place.

You may need to read it again.

What Minims Oxybuprocaine is used for

Minims Oxybuprocaine belongs to a group of medicines called local anaesthetics.

It is used to numb or block the feeling of pain in your eye(s) during short-term medical procedures. These procedures may include, measuring the pressure in your eye, inserting contact lenses, removing foreign bodies from your eye(s) and other minor operations on your eye(s).

Ask your doctor if you have any questions about why this medicine has been prescribed to you.

Your doctor may have prescribed it for another reason.

Before you are given Minims Oxybuprocaine

When you must not be given it

Do not use Minims Oxybuprocaine if you have an allergy to:

- Any medicine containing Oxybuprocaine
- Any other local anaesthetic
- Any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- Shortness of breath
- Wheezing or difficulty breathing
- Swelling of the face, lips, tongue or other parts of the body
- Rash, itching or hives on the skin

Do not use this medicine if you have an infection in your eye.

Do not use this medicine if you are pregnant or breast-feeding.

Your doctor will discuss the risks and benefits involved.

Do not use this medicine after the expiry date printed on the pack or

if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

This medicine has been developed for use in the eye only. It should not be swallowed or injected.

If you are not sure whether you should be given this medicine, talk to your doctor.

Before you are given it

Tell your doctor if you have had an allergy to any other medicines, foods, preservatives or dyes.

Tell your doctor if you are pregnant or breast-feeding.

Your doctor will discuss the risks and benefits involved.

If you have not told your doctor about any of the above, tell him/ her before you start using Minims Oxybuprocaine.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Minims Oxybuprocaine may interfere with each other. These include a group of medicines called anticholinesterases.

These medicines may be affected by Minims Oxybuprocaine or may affect how well it works. You may need different amounts of your medicines,

MINIMS® OXYBUPROCAINE

or you may need to use different medicines.

Your doctor will advise you and decide whether or not to give you Minims Oxybuprocaine.

How Minims Oxybuprocaine is given

Follow all directions given to you by your doctor carefully.

They may differ from the information contained in this leaflet.

How much to use

Your doctor will put the correct number of drops of Minims Oxybuprocaine into your eye(s) before starting the procedure on your eye(s). Your doctor will not give you a prescription for your own use. This medicine has been specifically developed for use by your doctor to numb the feeling in your eye(s) while he/she is attending to your eye(s).

The usual dose is one drop into your eye(s) before starting the procedure on your eye(s). The number of times your doctor will put drops in your eye(s) will depend on the type and length of the procedure to be performed. For most procedures one or two drops will be sufficient. For the removal of foreign bodies or minor surgery, three to six drops may be required during the procedure.

If you are wearing contact lenses, you will be asked to remove them before putting in the eye drops. You may put your contact lenses back into your eye after the numbness has worn off or when you doctor tells you to.

Your eye(s) will remain numb for up to one hour after the last drop is put in.

Do not touch or rub your eye while your eye is numb.

You may scratch the surface of the eye and damage it. Your doctor will put a patch over your eye after

finishing the procedure to protect your eye and stop you from touching it.

If you use too much (overdose)

The doctor giving you Minims Oxybuprocaine will be experienced in its use, so it is unlikely that you will be given an overdose.

If too many drops have accidentally been put into your eye(s), immediately rinse your eye(s) with warm water.

Accidental swallowing of the solution is unlikely to cause any bad side effects due to the low content of oxybuprocaine. If you think that you or anyone else may have swallowed any or all of the contents of the container of Minims Oxybuprocaine, immediately telephone your doctor or Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766) for advice.

After being given Minims Oxybuprocaine

Things you must do

Protect your eyes from dust.

You should protect your anesthetised eye from dust particles, which could cause infections.

Things you must not do

Do not touch or rub your eye while your eye is numb.

You may scratch the surface of the eye and damage it. Your doctor will put a patch over your eye after finishing the procedure to protect your eye and stop you from touching it.

Things to be careful of

Be careful driving or operating machinery until you know how

Minims Oxybuprocaine affects vou.

This medicine generally does not cause any problems with your ability to drive a car or operate machinery. However, it may cause transient blurred vision at the time of application and sufficient time should be allowed to regain clear vision before driving or operating hazardous machinery.

Side effects

Tell your doctor or pharmacist if you do not feel well while you are using Minims Oxybuprocaine.

This medicine helps most people to numb the eye, but it may have unwanted effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects.

You may not experience any of them.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- Stinging or burning sensation
- Prolonged blurred vision and / or problems seeing clearly
- Local irritation or feeling of having something in the eye
- Inflammation of the eye
- Skin rash
- Nausea and vomiting

The above list includes the more common side effects of your medicine. They are usually mild and short lived.

Tell your doctor immediately or visit Accident and Emergency at your nearest hospital if you notice any of the following:

 Swelling of the face, lips, mouth, tongue or throat or other parts of the body

- Slow heart beat
- Drowsiness, confusion, agitation, feeling of extreme happiness, disorientation, visual, hearing or speech disorders

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are rare.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice any other effects.

After using Minims Oxybuprocaine

Storage

Minims Oxybuprocaine will be stored by your doctor in a refrigerator (2-8°C). The eye drops should not be frozen or exposed to strong light.

Product description

What it looks like

Minims Oxybuprocaine comes in single dose units containing 0.5mL of a clear, colourless solution.

Each carton contains 20 single dose units.

Ingredients

Minims Oxybuprocaine contains 0.4% w/v of Oxybuprocaine as the active ingredient.

It also contains:

- · Water Purified
- Hydrochloric acid

The eye drops do not contain any preservatives.

Manufacturer/Supplier

In Australia, Minims Oxybuprocaine is supplied by:

Bausch & Lomb (Australia) Pty Ltd

Chatswood, NSW 2067

Phone: 1800 251 150 Minims

Oxybuprocaine 0.4%

AUST R 32259

In New Zealand, Minims
Oxybuprocaine is supplied by:

Bausch & Lomb (NZ) Ltd

c/- Bell Gully

Auckland Vero Centre

48 Shortland Street

Auckland 1140

New Zealand

Toll free number: 0508375394

Date of preparation

This leaflet was prepared in

February 2018